

New study quantifies natural flux of methane gas in the northeast Pacific

Submitted by Katie Shoemaker Tue, 2018-09-11 14:42

Beneath the ocean floor, bacteria produce methane gas that is regularly released up through the sediment and into the sea water as bubble streams (*Figure 1*). While these gas flares have been observed on continental margins around the world, until now there has been no systematic study of all available gas flow observation data to estimate the total amount of methane escaping from the seafloor. These data are important for the global inventory of carbon and also for analyzing the uptake of carbon dioxide (ocean acidification), and its impact on climate change.



Figure 1: Methane gas bubbles escaping the seafloor near Clayoquot Slope.

In a new study recently published in [Nature.com](https://www.nature.com), researchers in Germany and Canada have now analyzed all available historic and recent gas bubble data from the Cascadia margin off British Columbia, Washington and Oregon (Figure 2).




Figure 2: Regional map of the Cascadia margin showing flare distribution and gas bubbling flow rate. The size of the blue circles is proportional to the gas flow rate; black circles show

vent sites without flow estimates. (Credit: Nature Communications)

Lead by GEOMAR's Dr. Michael Riedel with his local colleagues in Kiel, along with researchers at MARUM Center for Marine Environmental Sciences and Department of Geosciences at University of Bremen, Ocean Networks Canada and the University of Victoria, the study used ship-based acoustic surveys of the water column, as well as near-seafloor acoustic and visual data. They gathered over 1100 vent sites with over 430 gas flow estimates for the entire Cascadia offshore margin.

ONC scientists Martin Scherwath and Martin Heesemann contributed to the research by providing ONC data—including ship-based acoustic data, underwater video and seafloor multi-beam sonar data—as well as helping to interpret and statistically summarize the data, and writing the manuscript.




Figure 3. Examples of acoustic data showing gas venting. (a) Gas flare in ~120 metres water depth from expedition FK009A (EK60), (b) example of flares in ~85 metres water depth from expedition MF0903 (EK60), (c) example of a flare in ~920 metres water depth from expedition ONC2014 (EK60), and (d) fan-view of EM302 data of flares in ~1335 metres water depth from expedition Naut2016. (Credit: Nature Communications)

?With so many ship surveys and relevant data sets readily available across the Cascadia margin (*Figure 3*), we just had to go ahead with the study and reveal where all the undiscovered vents were and how much the total gas flux was likely to be.? comments
ONC?s Martin Scherwath.

Since 2010, ONC has deployed multi-beam sonar instruments at Clayoquot Slope to study plumes of methane bubbles escaping from the seafloor. Previous research has monitored the bubble fluctuations at this site, but it is practically impossible to observe an entire margin's methane activity at all times. In order to obtain a gas flux estimate for the whole region, the existing observations were extrapolated in space and time to estimate gas activity over the entire margin, even at places where no data exist.

?While this estimate has large uncertainties,? says Martin Scherwath ?the research suggests that for every square metre of seafloor, about one gram of methane is bubbling up every year. This amounts to 100,000 tonnes of methane escaping the sediments into the ocean in offshore Cascadia every year.?

The existing observations at the Cascadia margin indicate that all of the gas is dissolving in the ocean, which means it is not escaping into the atmosphere where methane would act as a powerful greenhouse gas. Rather, its ocean dissolution is exacerbating ocean acidification.

[Read the full research paper on Nature.com](#)

Tags:

- [methane](#)
- [barkley canyon](#)
- [gas hydrates](#)

Categories:

- [News Stories](#)

```
// FIXES AMPERSAND IN BREADCRUMB var ONC_breadcrumb =  
document.getElementById("breadcrumb"); if (ONC_breadcrumb) { var ONC_innerHTML =  
ONC_breadcrumb.innerHTML; ONC_innerHTML = ONC_innerHTML.replace("&", "&");  
ONC_breadcrumb.innerHTML = ONC_innerHTML; }
```

Highlights

- [Audio](#)
- [Data](#)
- [Learning](#)

- [Science](#)
- [Video](#)

Reading Room

- [Active Research](#)
- [Backgrounders](#)
- [FAQs](#)
- [Glossary](#)
- [News Briefs](#)
- [News Stories](#)
- [Newsletters](#)
- [Publications](#)

Cool Stuff

- [Apps](#)
- [Digital Fishers](#)
- [iBooks & e-Pubs](#)
- [Live Video](#)
- [Maps](#)
- [Images](#)
- [State of the Ocean](#)

Data & Tools

- [Apps](#)
- [Data Plots](#)
- [Data Search](#)
- [Data Policy](#)
- [Data Help](#)
- [OPeNDAP Web Services](#)

Opportunities

- [Calendar](#)
- [Educator Opportunities](#)

- [Global Partnerships](#)
- [Industry Network](#)
- [Jobs](#)
- [Staff List](#)
- [Technology Services](#)

Sites & Instruments

- [Arctic Sites](#)
- [Northeast Pacific Sites](#)
- [Salish Sea Sites](#)
- [Notice to Mariners](#)

Follow Us



[Sign up for our newsletter](#)

Feedback

Send us your questions and comments *

How could we improve this page?

Your Name

Your Email *

Your Location

CAPTCHA

This question is for testing whether or not you are a human visitor and to prevent automated spam submissions.



What code is in the image? *
Enter the characters shown in the image.



[About Us](#) | [Contact Us](#) | [Media Relations](#) | [Legal Notices](#)

© Ocean Networks Canada. All rights reserved. 2474 Arbutus Road, Victoria, BC, V8N 1V8
| 1.250.472.5400

```
(function () { var d = new Date; var year = d.getFullYear();  
document.getElementById("copyright-date").innerHTML = year; })();
```

Source URL: <https://oceannetworks.ca/new-study-quantifies-natural-flux-methane-gas-northeast-pacific>